

Holographic Noise in Atomic Systems and Optical Cavities

C. Lämmerzahl and E. Göklü



Centre for Applied Space Technology and Microgravity (ZARM),
University of Bremen, 28359 Bremen, Germany

Holographic noise workshop
AEI, Hannover, 19. – 20.5.2009



1 Introduction

- 1 Introduction
- 2 Metric fluctuations in atomic systems
 - The basic equations
 - Equivalence principle
 - Decoherence
 - Spreading of wave packets

- 1 Introduction
- 2 Metric fluctuations in atomic systems
 - The basic equations
 - Equivalence principle
 - Decoherence
 - Spreading of wave packets
- 3 Metric fluctuations in optical cavities


- 1 Introduction
- 2 Metric fluctuations in atomic systems
 - The basic equations
 - Equivalence principle
 - Decoherence
 - Spreading of wave packets
- 3 Metric fluctuations in optical cavities
- 4 Summary and outlook

Outline

- 1 Introduction
- 2 Metric fluctuations in atomic systems
 - The basic equations
 - Equivalence principle
 - Decoherence
 - Spreading of wave packets
- 3 Metric fluctuations in optical cavities
- 4 Summary and outlook

General remarks I

two-step approach / situation



- **First step:** Derivation of metrical noise from fundamental principles
 - Existence of some metrical noise is generally accepted
 - May perhaps be more general (torsional noise, noise of non-metricity, Finslerian noise, ...)
- **Second step:** search for all consequences of metrical fluctuations (not only holographic noise)
 - search for most promising experiments
 - does not depend on how holographic noise is derived or founded


General remarks II

Geometric noise is **universal**

- geometric noise should appear in all physical systems
 - optical interferometers
 - optical cavities
 - atomic interferometer
 - atoms with spin
 - atomic, molecular energy levels
- all should be consistent with existing data / measurements
- should not depend on temperature, on charge, ...
- characteristics ?

It is not enough to see this noise in one system. **To experimentally show the universality is mandatory for the proof of its existence.**

Classes of experiments



Classes of experiments

Much has been done, but this needs more theory:

- calculation of all experimental consequences of space–time fluctuations
- then carrying through all promising experiments

It is important that this noise is **seen in more than one physical system**

Nothing has been seen until now, except perhaps in GEO600

Classes of experiments

Much has been done, but this needs more theory:

- calculation of all experimental consequences of space–time fluctuations
- then carrying through all promising experiments

It is important that this noise is **seen in more than one physical system**

Nothing has been seen until now, except perhaps in GEO600

Is part of the big task: Search for new physics

- violation of Universality of Free Fall
- violation of Universality of Gravitational Redshift
- violation of Lorentz Invariance
- modified gravitational field
- fundamental noise (provides **new window**)

Some history I

space–time foam, fuzzy space–time, space–time fluctuations
goes back to Hawking 1973, Bekenstein 1975, Wheeler 1982

- **Ellis, Hagelin, Nanopoulos & Srednicki 1984** decoherence due to fluctuating metric, neutron interferometry
- **Percival & Strunz 2000** Influence of stochastic metric fluctuations on atom interferometry
- **Power & Percival 2000** Decoherence of wave packets from conformal space–time fluctuations, modified Schrödinger equation for density matrix
- **Amelino–Camelia 2000** Saleker–Wigner argument, random–walk, modified dispersion

yields general Brownian motion ansatz $S_{\text{sf}}^{(\alpha, \gamma)}(\nu) = \zeta \frac{\Lambda}{c} \left(\frac{l_{\text{Planck}}}{\Lambda} \right)^\alpha \left(\frac{\nu}{c/\Lambda} \right)^\gamma$

- **Ng & van Dam 2000** Distance measurement by clocks (based on Saleker–Wigner argument) $\delta g \sim (l_{\text{Planck}}/l)^{\frac{2}{3}}$

Some history II

- **Ng 2002** Holographic principle $(l/\delta l)^3 \leq (l_{\text{Planck}}/l) \Rightarrow \delta g \sim (l_{\text{Planck}}/l)^{\frac{2}{3}}$.
Idea: holographic principle follows from space–time fluctuations
Relation to quantum computing
Influence on dispersion relations, decoherence of light phase, UHECR, non–locality, ...
- **Hu & Verdaguer 2002, 2008** Axiomatic approach: Einstein–Langevin equations, application to backreaction problems and black hole fluctuations
- **Ford 2003 – 2008** No specific model, luminosity fluctuations, line broadening, angular blurring, black hole fluctuations
- **Aloisio, Galante, Grillo, Liberati, Lucio & Mendez 2006** no specific scenario, relation to modified dispersion
- **Wang, Bonifacio, Bingham & Mendonca 2006, 2009** Conformal fluctuations and decoherence of quantum particle, effect for very large masses $\sim 10^{19}$ a.m.u.
- **Hogan 2008** Holographic noise in GEO600

This talk

Recent work at ZARM

Influence of space–time fluctuations on quantum systems

- fundamental noise in optical resonators, [Schiller, Lämmerzahl, Müller, Braxmaier, Herrmann & Peters, PRD 2004](#) (experiment)
- apparent violation of weak equivalence principle, [Göklü & Lämmerzahl, CQG 2008](#)
- decoherence, [Breuer, Göklü & Lämmerzahl, CQG 2008](#)
- spreading of wave packets, [Göklü, Lämmerzahl, Camacho & Macias, 2009](#)

our motivation ...

Bremen Drop Tower of ZARM



Tower 146 m

drop tube
110 m


free fall time
 $= 4.7 \text{ s}$

deceleration
 $\sim 30 \text{ g}$


Bremen Drop Tower of ZARM




BEC in microgravity



design of capsule




vacuum chamber



capsule

BEC in microgravity – long free evolution


50 ms



100 ms



500 ms




1000 ms

10^4 atoms, 1 s free evolution time (not possible on ground)

BEC in microgravity – long free evolution


50 ms




100 ms



500 ms



1000 ms



10^4 atoms, 1 s free evolution time (not possible on ground)

Outline

- 1 Introduction
- 2 Metric fluctuations in atomic systems
 - The basic equations
 - Equivalence principle
 - Decoherence
 - Spreading of wave packets
- 3 Metric fluctuations in optical cavities
- 4 Summary and outlook

Outline

- 1 Introduction
- 2 Metric fluctuations in atomic systems
 - The basic equations
 - Equivalence principle
 - Decoherence
 - Spreading of wave packets
- 3 Metric fluctuations in in optical cavities
- 4 Summary and outlook

The basic equations

The model

- Klein–Gordon equation

$$g^{\mu\nu} D_\mu D_\nu \varphi + m^2 \varphi = 0, \quad D = \partial + \{ \cdot \}$$

- Fluctuating metric

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}, \quad |h_{\mu\nu}| \ll 1$$

- noise

$$\langle h_{\mu\nu}(x) \rangle_{\text{st}} = \gamma_{\mu\nu}, \quad \delta^{\rho\sigma} \langle h_{\mu\rho}(x) h_{\nu\sigma}(x) \rangle_{\text{st}} = \sigma_{\mu\nu}^2$$

- small amplitude of fluctuations
- frequency might be large
- wavelength might be small
- we do not require the $h_{\mu\nu}$ to obey a wave equation

The basic equations

Approximations

- Weak field up to second order $\tilde{h}^{\mu\nu} = h^{\mu\rho} h_{\rho}^{\nu}$
- Relativistic approximation of metric and quantum field (à la Kiefer and Singh)

$$\begin{aligned}
 H\psi = & -({}^{(3)}g)^{\frac{1}{4}} \frac{\hbar^2}{2m} \Delta_{\text{cov}} \left(({}^{(3)}g)^{-\frac{1}{4}} \psi' \right) + \frac{m}{2} \left(\tilde{h}_{(0)}^{00} - h_{(0)}^{00} \right) \psi \\
 & - \frac{1}{2} \left\{ i\hbar \partial_i, h_{(1)}^{i0} - \tilde{h}_{(1)}^{i0} \right\} \psi
 \end{aligned}$$

manifest hermitean w.r.t. flat scalar product

- only second order terms do not vanish by averaging

Outline

- 1 Introduction
- 2 Metric fluctuations in atomic systems
 - The basic equations
 - **Equivalence principle**
 - Decoherence
 - Spreading of wave packets
- 3 Metric fluctuations in optical cavities
- 4 Summary and outlook

Short wavelength

Spatial average

- spatial average

$$\langle A \rangle_s(x) := \frac{1}{V_x} \int_V A(y) d^3y$$

- short wavelength of fluctuations: V small $\Rightarrow \langle \psi \rangle_s(x) = \psi(x)$
- spatial average of Schrödinger equation

$$H = \frac{1}{2m} (\delta^{ij} + \alpha^{ij}(x)) p_i p_j + \alpha_0$$

with $\alpha(x) = \langle \tilde{h}^{ij} - h^{ij} \rangle_s(x)$

- $\alpha^{ij}(x)$: small variation w.r.t. x , fluctuations w.r.t. t .
- decompose $\alpha^{ij}(x) = \tilde{\alpha}^{ij}(x) + \gamma^{ij}(x)$ with $\langle \gamma^{ij} \rangle_t = 0$
- $\tilde{\alpha}^{ij}(x)$ acts like an anomalous inertial mass tensor

Space-time fluctuations

Fluctuation model

- $\alpha^{ij} \leftrightarrow$ spectral noise density of fluctuations
- particular model:

$$\tilde{\alpha}^{ij}(x) = \frac{1}{V_x} \int_{V_x} \tilde{h}^{ij}(x, t) d^3x = \frac{1}{V_x} \int_{1/V_x} (S^2(k, t))^{ij} d^3k$$

- model: power law spectral noise density

$$(S^2(k, t))^{ij} = (S_{0n}^2)^{ij} |k|^n \quad \Rightarrow \quad \alpha^{ij}(x) = (S_{0n}^2)^{ij} \lambda_p^{-(6+n)}$$

with $\dim(S_{0n}^2)^{ij} = \text{length}^{3+\frac{n}{2}}$

- $\lambda_p =$ length scale of particle $= \lambda_{\text{Compton}}$
- $V_x \sim \lambda_p^3$

Space-time fluctuations

Fluctuation model

- assumption: $S_{0n} \sim l_{\text{Planck}}^{3+\frac{n}{2}}$, then

$$\alpha^{ij}(x) \sim \left(\frac{l_{\text{Planck}}}{l_{\text{Compton}}} \right)^{\beta} a^{ij}(x), \quad \beta = 6 + n, \quad a^{ij}(x) = \mathcal{O}(1)$$

- effective Hamiltonian

$$H = \frac{1}{2m} \left(\delta^{ij} + \left(\frac{l_{\text{Planck}}}{l_{\text{Compton}}} \right)^{\beta} a^{ij}(x) \right) p_i p_j = \frac{1}{2m} \left(\delta^{ij} + \frac{\delta m^{ij}(x)}{m} \right) p_i p_j$$

δm^{ij} = anomalous inertial mass tensor, depends on particle

- δm^{ij} leads to violation of Universality of Free Fall ([Haugan 1979](#))

- $\beta = \frac{1}{2} \leftrightarrow$ random walk
- $\beta = \frac{2}{3} \leftrightarrow$ holographic noise

Result

Result

metric fluctuations \Rightarrow anomalous inertial mass \rightarrow **apparent** violation of UFF

- alternative route for violation of UFF and LLI

Example

for Cesium and Hydrogen:

$$\eta_{\beta=1} = 10^{-17}, \quad \eta_{\beta=2/3} = 10^{-12}, \quad \eta_{\beta=1/2} = 10^{-9}$$

$\beta = \frac{1}{2}$ already ruled out.

(Göklü & C.L. CQG 2008)

are there transverse effects?

Outline

- 1 Introduction
- 2 Metric fluctuations in atomic systems
 - The basic equations
 - Equivalence principle
 - **Decoherence**
 - Spreading of wave packets
- 3 Metric fluctuations in optical cavities
- 4 Summary and outlook

Decoherence

The model

- model as above

$$H = \frac{1}{2m} (\delta^{ij} + \tilde{\alpha}^{ij} + \gamma^{ij}(t)) p_i p_j$$

- discuss now the influence of γ^{ij} and neglect $\tilde{\alpha}^{ij}$
- neglect small x -dependence

Noise model

- isotropic fluctuations $\gamma^{ij}(t) = \sigma \delta^{ij} \xi(t)$
 - white noise $\langle \xi(t) \rangle = 0$, $\langle \xi(t) \xi(t') \rangle = \delta(t - t')$
 - $\dim \sigma = \text{time} = \tau_c$
-
- practically no influence from colored noise
 - $\gamma^{ij}(t)$ random process

Decoherence

Master equation

- stochastic Schrödinger equation in interaction picture

$$i\hbar \frac{d}{dt} |\tilde{\psi}\rangle = \tilde{H}_\gamma |\tilde{\psi}\rangle, \quad \tilde{H} = e^{\frac{i}{\hbar} H_0 t} H_\gamma e^{-\frac{i}{\hbar} H_0 t}$$

with random Hamiltonian \tilde{H}_γ with $\langle \tilde{H}_\gamma \rangle_t = 0$

- averaging over fluctuations \Rightarrow averaged density matrix

$$\tilde{\rho}(t) = \langle |\tilde{\psi}\rangle \langle \tilde{\psi}| \rangle$$

- master equation for averaged density matrix to second order in the fluctuations

$$i\hbar \frac{d}{dt} \tilde{\rho} = -\frac{i}{\hbar} \int_0^t \langle [\tilde{H}_\gamma(t), [\tilde{H}_\gamma(t'), \tilde{\rho}(t)]] \rangle dt'$$

Decoherence

Markovian master equation

- in Schrödinger picture

$$i\hbar \frac{d}{dt} \rho(t) = [H_0, \rho(t)] + i\hbar (\mathcal{D}\rho)(t)$$

with

$$(\mathcal{D}\rho)(t) = -\frac{1}{2}[V, [V, \rho(t)]] \quad \text{with} \quad V = \frac{\sqrt{\tau_c}}{\hbar} \frac{p^2}{2m}$$

- master equation is in Lindblad form \Rightarrow defines a completely positive quantum-dynamical semigroup
- energy is conserved
- \mathcal{D} is the dissipator

Decoherence

Decoherence time

- solution of master equation in momentum space

$$\rho(p, p', t) = \exp \left(-\frac{i}{\hbar} \Delta E t - \frac{(\Delta E)^2 \tau_c}{2\hbar^2} t \right) \rho(p, p', 0)$$

decoherence time

$$\tau_D = \frac{2\hbar^2}{(\Delta E)^2 \tau_c} = 2 \left(\frac{\hbar}{\Delta E \tau_c} \right)^2 \tau_c$$

- for $\tau_c = t_{\text{Planck}}$

$$\tau_D = \frac{10^{13} \text{ s}}{(\Delta E / \text{eV})^2}$$

- too large for being observable
- may change for BECs

(Breuer, Göklü & C.L. 2009)



Outline

- 1 Introduction
- 2 Metric fluctuations in atomic systems
 - The basic equations
 - Equivalence principle
 - Decoherence
 - Spreading of wave packets
- 3 Metric fluctuations in optical cavities
- 4 Summary and outlook

Spreading of wave packets

Model

- dynamics: same model as above

$$H = H_0 + V(x), \quad V(x) = \mathcal{O}(\hbar \partial \partial \hbar, \partial \hbar \partial \hbar)$$

- V is Gaussian random function

$$\langle V(x) \rangle = 0, \quad \langle V(x), V(x') \rangle = V_0^2 \delta(t - t') g(x - x')$$

longer calculations ...

The spreading

for Gaussian correlation and Gaussian initial wave packet


$$\langle x^2(t) \rangle = \underbrace{\sigma^2 + \frac{\hbar^2}{4m^2\sigma^2}t^2}_{\text{free evolution}} + \underbrace{\frac{\sigma_{px}}{m}t}_{\text{diffusion}} + \underbrace{\frac{V_0}{3\sqrt{2\pi}m^2a^3}t^3}_{\text{superdiffusion}} \quad (1)$$

(Göklü, C.L., Camacho & Macias, in preparation)


Outline

- 1 Introduction
- 2 Metric fluctuations in atomic systems
 - The basic equations
 - Equivalence principle
 - Decoherence
 - Spreading of wave packets
- 3 Metric fluctuations in in optical cavities**
- 4 Summary and outlook


Optical resonators



Optical resonators



Optical resonators



M. Tobar
ZARM

Metric fluctuations in cavities

Ansatz for the strain noise spectrum (Amelino–Camelia PRD 2000)

$$\left(\frac{\Delta L}{L}\right)^2 = \int S_{\text{sf}}(\nu) d\nu \quad \text{with} \quad S_{\text{sf}}(\nu) = \zeta \frac{\Lambda}{c} \left(\frac{L_{\text{Pl}}}{\Lambda}\right)^\beta \left(\frac{\nu}{c/\Lambda}\right)^\gamma,$$

Λ = length characteristic of experimental setup

Specification for two random-walk hypotheses:

$$S_{\text{sf}}^{\beta=1, \gamma=-2} = 5 \cdot 10^{-27} \zeta_{\text{rw1}} \left(\frac{\text{m}}{\Lambda}\right)^2 \left(\frac{\text{Hz}}{\nu}\right)^2 \text{Hz}^{-1}$$

$$S_{\text{sf}}^{\beta=2, \gamma=-2} = 7 \cdot 10^{-62} \zeta_{\text{rw2}} \left(\frac{\text{m}}{\Lambda}\right)^3 \left(\frac{\text{Hz}}{\nu}\right)^2 \text{Hz}^{-1},$$

Devices

- low frequencies and small devices are preferred setups
- **Optical resonators**: access to μHz range.

data of a frequency comparison between two optical resonators has been analyzed



Metric fluctuations in cavities

measured quantity = $\nu_2 - \nu_1$

$$\begin{aligned} \delta \left(\frac{\nu_2 - \nu_1}{\nu_1} \right) &= \left(\frac{\delta L_2^{\text{sf}}}{L_2} - \frac{\delta L_1^{\text{sf}}}{L_1} \right) \\ &+ \left(\frac{\delta L_2^{\text{phys}}}{L_2} - \frac{\delta L_1^{\text{phys}}}{L_1} \right) \\ &+ \left(\frac{\delta L_2^{\text{lock}}}{L_2} - \frac{\delta L_1^{\text{lock}}}{L_1} \right) + \dots \\ &= S_{\text{sf}} + S_{\text{phys}} + S_{\text{lock}} + \dots \end{aligned}$$


half of total noise represents upper bound to S_{sf} .

Comparison with RW ansätze

$$\zeta_{\text{rw1}} \leq 2 \cdot 10^{-13} \quad \zeta_{\text{rw2}} \leq 4 \cdot 10^{20}.$$

Parameters are of order 1: rules out RW hypothesis 1 ([Schiller et al PRD 2004](#)).

For identification of effect one has to use many materials



setup A: resonators parallel in different cryostats

setup B: two cavities orthogonally in same cryostat


Holographic noise in cavities

Holographic noise

- Spectral noise density for holographic scenario

$$S_{\text{sf}}(\nu) = \zeta \frac{\Lambda}{c} \left(\frac{L_{\text{Pl}}}{\Lambda} \right)^{\beta} \quad \beta = \frac{2}{3}$$

- for cavity $S_{\text{sf}}(\nu) \sim 10^{-33} \text{ Hz}^{-1}$



Outline

- 1 Introduction
- 2 Metric fluctuations in atomic systems
 - The basic equations
 - Equivalence principle
 - Decoherence
 - Spreading of wave packets
- 3 Metric fluctuations in in optical cavities
- 4 Summary and outlook

Summary and outlook

Summary

Metric (holographic) noise appears as

- apparent violation of weak equivalence principle
- decoherence
- additional spreading of wave packets

Required

- to proof universality
- to discrimination with other noise surces
- to show cosistency

Outlook

- description of more experiments (effect on spin or helicity, ...)
- effects on BECs (long evolution time – free fall of BECs)

Spin and space-time fluctuations (roughness)

- Geodesic equation for static spherically symmetric metric

$$\left(\frac{dr}{ds}\right)^2 = \frac{1}{g_{tt}g_{rr}} \left(E^2 - g_{tt} \left(\epsilon + m \frac{L^2}{r^2} \right) \right)$$

- Assuming $g_{rr} = 1/g_{tt}$ and $g_{tt} = 1 + h \cos(kr)$ (everywhere C^∞)
Then for radial motion ($L = 0$)


$$\left(\frac{dr}{ds}\right)^2 = E^2 - 1 - h \cos(kr)$$

Can be solved by elliptic function

- for $h \ll 1$

$$r = \sqrt{E^2 - 1} s + h \frac{\sin(\sqrt{E^2 - 1} k s)}{2(E^2 - 1)k}$$

like zitterbewegung



Spin and space-time fluctuations

Kretschmann scalar

$$K = R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} \sim \frac{g''_{tt}}{g_{tt}} \sim hk^2 \frac{\cos(kr)}{1 + h \cos(kr)}$$

Consequence

- For $k \rightarrow \infty$: solution of geodesic equation approaches straight line
 - Kretschmann scalar $\rightarrow \infty$
- \Rightarrow point particles do not see fluctuating curvature
 particles with spin are sensitive to curvature $a = R(\cdot, u, S, u)$
- may be of importance in atomic interferometry, spectroscopy, ...
 - if fluctuations are of quantum gravity origin \rightarrow estimates
 - needs to be compared with analysis of Dirac equation in fluctuating space-time metric

Göklü & C.L. in preparation

One final remark

- I think one should look for experiments with other physical systems, e.g., atoms, BECs, ...
- Then one has more possibilities to check consistency with existing data for the other experiments

Thank you !

Thanks to

- German Reserach Foundation DFG
- Center of Excellence QUEST
- German Aerospace Agency DLR